

The Impact and Innovative Remediation Technologies of Hazardous Waste Disposal on Land

By

^aInnocent Ojeba Musa, ^aMustapha Abdulsalam, ^bMiracle Uwa Livinus, and
^cOluwafemi Adebayo Oyewole*

Abstract

Land disposal of hazardous waste is one of the worst issues with regard to environmental management. Amid the rising wave of industrial, agricultural, and biomedical discoveries, the challenge of disposing of waste through land has become a widespread menace to soil fertility, balance in the ecosystem, and health. The present study would look at the various impacts that hazardous waste has on both the land system and the soil system and would then study in detail the advanced methods of the treatment of hazardous waste, like the technology of nanotechnology treatment, electro-treatment, and genetic engineering microbial intervention. We believe that enduring legacies of hazardous waste can only be tackled by an integrated framework that is a blend of environmental thinking, coupled with technological inventiveness. The current study provides a review of the existing knowledge base, introduces the new developments, and provides a future perspective on land stewardship.

Keywords: Hazardous waste, land pollution, remediation, nanotechnology, phytoremediation, electrokinetic remediation, environmental justice.

1. Introduction

Soil is usually undervalued in its real value, even though it is the basis of earthly life. It is not simply soil under our feet, but the biological soil is a working environment that is comprised of minerals, organic matter, air, water, and a rich population of microorganisms. Soil harbors biodiversity facilitates food production, filters water, and aids in the regulation of the carbon cycle on the planet. However, this invaluable commodity is in great danger, much of which is as a result of the irresponsible disposal of toxic waste¹. Although the planet is capable of cleaning itself to some extent, it cannot cope with uncontrollable amounts of pollution produced by mankind. The amount of hazardous waste generated in the world each year, which reaches above 400 million tonnes, is mishandled most of the time, with mismanagement typically involving throwing it into open land or landfills that are mostly poorly controlled². Unrestrained urbanization, industrial activity combined with oversaturating of farming with chemicals, has only fueled the problem

^a Department of Microbiology, Skyline University, Kano, Kano State, Nigeria

^b Department of Biochemistry, Skyline University, Kano, Kano State, Nigeria

^c Department of Microbiology, Federal University of Technology Minna, Niger State, Nigeria

* Corresponding Author: Email address: oa.oyewole@futminna.edu.ng; Tel: +234-8140031875

¹ Heacock, M., Kelly, C. B., Asante, K. A., Birnbaum, L. S., Bergman, Å. L., Bruné, M. N., Buka, I., Carpenter, D. O., Chen, A., Huo, X., Kamel, M., Landrigan, P. J., Magalini, F., Diaz-Barriga, F., Neira, M., Omar, M., Pascale, A., Ruchirawat, M., Sly, L., Sly, P. D., ... Suk, W. A. (2016). E-Waste and Harm to Vulnerable Populations: A Growing Global Problem. *Environmental Health Perspectives*, 124(5), 550–555.
<https://doi.org/10.1289/ehp.1509699>

² Heacock .M. et al. (2016). Ibid.

further, resulting in a toxic cocktail of harmful chemicals such as heavy metals, petroleum products, and contaminated soils.

What is especially dangerous about hazardous waste is that it is persistent and toxic. Lead, cadmium, mercury, and polycyclic aromatic hydrocarbons (PAHs) are examples of substances that do not degrade readily. These wastes may remain in the soil for many years, unlike organic waste, which may decompose over a period of time, thereby destroying the physical, chemical, and biological balance of the soil. They damage the delicate microbial populations that are necessary for the healthy soil and are used in plant growth ³. Lower-income areas with poor waste management systems allow waste to be discarded in farmland, abandoned industrial areas, and areas where marginalized communities live without the funds to defend themselves. It leads to the long-term experience of poisoned air, water, and soil ⁴.

Heavy metals and chemicals in petroleum products are some of the most significant pollutants because they disturb key microbial cycles that recycle nutrients and sustain plant productivity. To take an example, cadmium and lead interfere with the symbiotic relationship between legumes and Rhizobium bacteria and decrease the fixation of nitrogen in the soil ⁵. In a related move, petroleum-based PAHs not only reduce soil fertility but also lead to severe health risks as they may leak into crops and groundwater, as well as cause cancer and genetic mutations ⁶.

In addition to these biological effects, hazardous waste is able to change the soil physically. Such wastes as fly ash in incinerators or coal combustion, or municipal waste, may alter the soil texture and water and nutrient holding capacity. Biomedical waste ash, usually containing high levels of toxic metals such as chromium, cadmium, and lead, may substantially change the pH of the soil and its floristic potential ⁷. Such consequences are not just superficial; the contaminants usually find their way into ground waters, which provide a very significant source of drinking water to billions of people. These toxins can potentially be accumulated in the human food chain because after their intake into the plants, they end up in the human food chain, leading to bioaccumulation and its long-term consequences⁸.

³Dubey, A., Malla, M.A., Khan, F., Chowdhary, K., Yadav, S., Sharma, S., Khare, P.K., Khan, M.L. (2019). Soil microbiome: a key player for conservation of soil health under a changing climate. *Biodiversity and Conservation*, 28:2405–2429.

⁴Livinus, M.U., Bala, S.Z., Abdulsalam, M., Innocent, M.O., Hassan, M. & Kini, P. (2024). Natural Occurrences of Soil Dilapidation. In: Prospects for Soil Regeneration and Its Impact on Environmental Protection. Cham: Springer Nature Switzerland; P. 205–223.

⁵Jach, M.E., Sajnaga, E. & Ziaja, M. (2022). Utilization of legume-nodule bacterial symbiosis in phytoremediation of heavy metal-contaminated soils. *Biology*, 11(5):676.

⁶Saheed, Y.K., Balogun, B.F., Odunayo, B.J. & Abdulsalam, M. (2023). Microarray gene expression data classification via Wilcoxon sign rank sum and novel Grey Wolf optimized ensemble learning models. *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, 20(6):3575–87.

⁷Alengebawy, A., Abdelkhalek, S. T., Qureshi, S. R., & Wang, M. Q. (2021). Heavy Metals and Pesticides Toxicity in Agricultural Soil and Plants: Ecological Risks and Human Health Implications. *Toxics*, 9(3), 42. <https://doi.org/10.3390/toxics9030042>

⁸Sonone, S.S., Jadhav, S., Sankhla, M.S. & Kumar, R. (2020). Water contamination by heavy metals and their toxic effect on aquaculture and human health through the food chain. *Letters in Applied NanoBioScience*; 10(2):2148–66.

But in reality, the menace of hazardous waste is much more than a mere environmental nuisance, and it is nothing less than a systemic crisis. It compromises food security, jeopardizes ecosystems, jeopardizes sustainable agriculture, and imposes a huge economic burden through loss of productivity and increased healthcare expenditures. Another important ethical concern is also voiced by it: are we leaving a poisonous legacy to the new generations.

Hazardous waste is an inevitable by-product of modernity rather than as a characteristic problem of the planetary future. This article addresses the issue of toxic pollution ruining soil systems and points out emergent remediation technologies that offer hope to repair the damage. What is more important, it requires us to change our mindset. It is high time to not only perform the cleanup of the polluted ground, but to reconsider the relation to the soil as to the very ground supporting life.

2. Environmental Significance and Societal Relevance

The technical dimension of hazardous waste management is just the tip of the iceberg, as hazardous waste management is also a matter of both social and environmental justice that spans across generations. Though the effect is critical, the problem is normally underrepresented in international policymaking. However, it is squarely related to a number of United Nations Sustainable Development Goals (SDGs) such as SDG 3 (Good Health and Well-being), SDG 6 (Clean Water and Sanitation), and SDG 15 (Life on Land). Such links demonstrate that not only does poor management of hazardous waste negatively affect the environment, but it also negatively impacts human health and frustrates any attempts at pursuing sustainable development.

One of the issues of concern here is the increased inequality in the disposal of waste. Most low and middle-income nations, especially in Sub-Saharan Africa, tend to discharge hazardous waste to open sites that are unregulated. Such dumpsites are often located close to marginalized societies and have no safety rules or supervision⁹. This is not just a coincidence; it is also a way of systemic negligence and exclusion of such communities in the decision-making process that directly impacts their environment and welfare¹⁰.

Consequently, these regions turn into what can be characterized by some scholars as the so-called toxic landscapes, the locations where industrial and medical waste is thrown away irresponsibly, perhaps even in the vicinity of crops. The existing communities in such places are condemned to use polluted air, water, and soil, and they have to live with the health and environmental costs of a problem they could hardly say anything about¹¹. Heavy metals and dioxins are openly burned together with electronic waste and industrial waste products that infiltrate the nearby waters and soil¹². Populations that have close exposure to informal biomedical waste pits are often

⁹ Yadav, H., Shah, D., Sayed, S., Horton, S. & Schroeder, L.F. (2021). Availability of essential diagnostics in ten low-income and middle-income countries: results from national health facility surveys. *Lancet Global Health* 9(11):e1553–60.

¹⁰ Kamoto, J., Clarkson, G., Dorward, P. & Shepherd, D. (2013). Doing more harm than good? Community-based natural resource management and the neglect of local institutions in policy development. *Land Use Policy*, 35:293–301.

¹¹ Kamoto, J. et al. (2013). Ibid.

¹² Grant, R., Oteng-Ababio, M. & Shin, M. (2024). Academic urban legend, Agbogbloshie: Sweeping away the “World’s Largest E-Waste Dumpsite”. *Habitat International*, 149:103097.

characterized by high morbidity of respiratory diseases, skin infections, and intestinal diseases ¹³. These are just a few real-life cases that are a graphic illustration of the consequences of irresponsible waste management of hazardous waste, which requires immediate and universal action.

Table 1: Common health impacts associated with land-based hazardous waste exposure, supported by epidemiological evidence.

Contaminant	Primary Exposure Route	Health Outcomes	Source Example
Lead	Ingestion/inhalation	Cognitive impairment, hypertension	Battery recycling sites (Ramírez et al. 2021)
Cadmium	Ingestion	Kidney dysfunction, bone demineralization	Industrial effluents (Lei et al. 2024)
Dioxins/PCBs	Ingestion (fatty foods)	Endocrine disruption, immune suppression	Open burning of e-waste (Fiolet et al. 2022)
PAHs	Inhalation/dermal contact	Carcinogenesis, reproductive toxicity	Petroleum spills (Montano et al 2025)
Biomedical waste (pathogens, mercury)	Airborne, contact	Zoonoses, antibiotic resistance, and heavy metal toxicity	Unregulated clinics (Kukreti et al. 2024)

Hazardous waste is not only toxic. Climate change is a multiplier since it can aggravate the mobility and toxicity of contaminants. The increase in temperature may increase the volatilization process of volatile organic compounds (VOCs), and unpredictable rainfall may facilitate the percolation of leachates into groundwater ¹⁴. It forms a vicious cycle where poor soil quality leads to impure agricultural products, which further decreases food security, causing rural citizens to move to cities, where many of them live in unofficial housing that has already suffered environmental hazards. Mathematically, the relationship between precipitation (P), contaminant concentration (C), and leachate migration depth (D) in unsaturated soils can be approximated using a modified advection–dispersion model:

$$D = \frac{qC}{\theta\lambda} \tag{1}$$

Where:

- q is the infiltration rate (cm/day),

¹³ Ravindra, K., Sareen, A., Dogra, S. & Mor, S. (2023). Appraisal of biomedical waste management practice in India and associated human health and environmental risk. *Journal of Environmental Biology*, 44(4):541–551.
¹⁴ Alao JO. (2025). The factors influencing the landfill leachate plume contaminants in soils, surface and groundwater, and associated health risks: A geophysical and geochemical view. *Public Health Environ.*1(1):20–43.

- C is the initial concentration of the contaminant (mg/L),
- θ is the soil moisture content,
- λ is the attenuation coefficient.

This equation illustrates how even modest increases in rainfall (q) can significantly deepen the vertical migration of hazardous substances, particularly in sandy or poorly buffered soils. As a result, what begins as a localized contamination event can quickly become a diffuse, landscape-scale problem.

From a societal perspective, these dynamics produce unequal burdens. Children in proximity to waste dumps are especially vulnerable due to their developing physiology and behaviors such as hand-to-mouth activity. Women involved in informal waste picking face higher risks of reproductive health problems due to chronic exposure to metals and biohazards¹⁵. Moreover, the economic impacts are substantial. Contaminated lands lose agricultural value, and healthcare systems are strained by preventable diseases linked to environmental exposures¹⁶. Policy inaction is partly a reflection of the compartmentalized nature of governance. Waste management is often treated in silos—technically as an engineering problem, rather than as a cross-sectoral challenge that touches on health, environment, justice, and development. However, integrated approaches such as the "One Health" model—which emphasizes the interconnectedness of human, animal, and environmental health—offer promising frameworks for reimagining waste governance¹⁷.

3. Nature and Mechanisms of Land-based Hazardous Waste Contamination

The terrestrial environment today is an unintended reservoir for a spectrum of hazardous materials originating from diverse anthropogenic activities. From the industrial heartlands of the global north to the informal waste zones of the global south, soil systems have absorbed decades of toxic legacies. The nature of these contaminants is as varied as their sources, each with distinct physicochemical properties and mechanisms of persistence that render remediation efforts technically challenging and resource intensive.

3.1 Typologies of Land-based Hazardous Waste

Hazardous waste disposed of on land can be categorized broadly into inorganic and organic pollutants, along with biological and radioactive constituents. Table 2 summarizes the major types, their origins, environmental behavior, and health risks:

¹⁵ Abdulsalam, M., Elelu, S.A., Innocent, M.O., Livinus, M.U., Latifat, S.O. & Mohammed, A.S. (2024). Women empowerment in environmental conservation. In: Prospects for soil regeneration and its impact on environmental protection. Cham: Springer. P. 351–370. Doi:10.1007/978-3-031-53270-2_7. ISBN: 978-3-031-53270-2.

¹⁶ Ramankutty, N., Mehrabi, Z., Waha, K., Jarvis, L., Kremen, C., Herrero, M., & Rieseberg, L. H. (2018). Trends in Global Agricultural Land Use: Implications for Environmental Health and Food Security. *Annual Review Of Plant Biology*, 69, 789–815. <https://doi.org/10.1146/annurev-arplant-042817-040256>

¹⁷ Mumford, E. L., Martinez, D. J., Tyance-Hassell, K., Cook, A., Hansen, G. R., Labonté, R., Mazet, J. A. K., Mumford, E. C., Rizzo, D. M., Togami, E., Vreedzaam, A., & Parrish-Sprowl, J. (2023). Evolution and expansion of the One Health approach to promote sustainable and resilient health and well-being: A call to action. *Frontiers in Public Health*, 10, 1056459. <https://doi.org/10.3389/fpubh.2022.1056459>

Table 2. Major categories of land-based hazardous wastes, their sources, environmental fate, and health effects.

Contaminant Type	Key Examples	Primary Sources	Environmental Behavior	Health Impacts
Heavy Metals	Lead (Pb), Cadmium (Cd), Mercury (Hg)	Battery recycling, mining, pigments	Binds to clay/humic particles; long-term persistence	Neurotoxicity, renal failure, carcinogenesis
Hydrocarbons	Benzene, PAHs, Toluene	Oil spills, fuel leakage, and refineries	Alters soil porosity; reduces water retention	Immunotoxicity, reproductive disorders
Persistent Organic Pollutants (POPs)	Dioxins, PCBs	E-waste burning, pesticides, and industrial lubricants	Lipophilic; bioaccumulate in biota	Endocrine disruption, developmental toxicity
Radionuclides	Cesium-137, Uranium-238	Nuclear accidents, medical isotopes	Alters soil pH and mineralogy	Genetic mutations, hematological malignancies
Biomedical Waste	PPE, syringes, pharmaceuticals	Hospitals, laboratories (especially post-COVID-19)	Infectious, chemically resistant	Pathogen transmission, antimicrobial resistance

3.2 Physicochemical and Biological Mechanisms of Soil Contamination

Hazardous wastes deposited in land combine with soil in very complex, and sometimes incompletely understood, combinations. The resulting contamination is affected by many factors that determine the degree and period of its effects, such as the chemical composition of waste, the physical and chemical composition of the soil, the activity of microorganisms, and even weather conditions during disposal¹⁸.

3.2.1 Heavy metals

Heavy metals, among other common soil pollutants, have been greatly studied- mainly because they are very toxic and are chemically stable. Organic matter and clay particles in the soil are prone

¹⁸ Meena, M. D., Yadav, R. K., Narjary, B., Yadav, G., Jat, H. S., Sheoran, P., Meena, M. K., Antil, R. S., Meena, B. L., Singh, H. V., Singh Meena, V., Rai, P. K., Ghosh, A., & Moharana, P. C. (2019). Municipal solid waste (MSW): Strategies to improve salt affected soil sustainability: A review. *Waste management* (New York, N.Y.), 84, 38–53. <https://doi.org/10.1016/j.wasman.2018.11.020>

to a strong binding with these metals. This does not make them less prone to leach into groundwater water, but it can be taken by the plants and microbes, which are pretty dangerous to the environment. As an illustration, cadmium has been found to disturb the natural nitrogen cycle by affecting the mutual beneficial association between legumes and Rhizobium bacteria and is able to affect the capacity of the soil to fix nitrogen biologically¹⁹. In the same way, mercury and lead have been associated with destabilization of the important enzymatic functions that weaken soil respiration and nutrient recycling.

3.2.2 Petroleum Derivatives and Hydrocarbons

Polycyclic aromatic hydrocarbons (PAHs), a common group of petroleum-based contaminants, can drastically alter soil structure. These compounds are lipophilic, meaning they tend to bind with fats and oils, which makes the soil more water-repellent. As a result, water penetration is reduced, harming plant root development²⁰. Additionally, these hydrocarbons can inhibit the growth of essential mycorrhizal fungi—organisms that play a critical role in helping plants absorb nutrients like phosphorus. The presence of these pollutants, therefore, not only changes the physical nature of the soil but also disrupts the biological relationships that keep it fertile and productive.

3.2.3 Persistent Organic Pollutants (POPs)

Chemicals such as polychlorinated biphenyls (PCBs) and dioxins are not easily degraded by photolysis, microorganisms, or chemicals. Such lipophilic compounds build up in fatty tissues and biomagnify through the trophic level. They are frequently found in the soil because of inappropriate electron waste disposal and pesticide contamination, and they are also associated with endocrine disruption and malformation of fetuses in mammals²¹.

3.2.4 Radionuclides

Long-term radiological dangers are caused by radionuclides like uranium and cesium, which are usually put there through nuclear accidents or improperly disposed of medical waste. They may replace such vital elements of the plant absorption plan as potassium and calcium, which entails radiation-induced hereditary damage to plants and soil inhabitants²². However, their decay over time alters the soil's mineralogical profile and induces oxidative stress in microbial cells.

¹⁹ Wu J.Y., & Yan S.D., 2024, Natural nitrogen boosters: the symbiotic relationship between legumes and rhizobia, *Molecular Soil Biology*, 15(2): 74-86. doi: 10.5376/msb.2024.15.0009.

²⁰ Shah, A. N., Tanveer, M., Shahzad, B., Yang, G., Fahad, S., Ali, S., Bukhari, M. A., Tung, S. A., Hafeez, A., & Souliyanonh, B. (2017). Soil compaction effects on soil health and crop productivity: an overview. *Environmental Science and Pollution Research International*, 24(11), 10056–10067. <https://doi.org/10.1007/s11356-017-8421-y>

²¹ Ghosh, A., Manna, M.C., Jha, S., Singh, A.K., Misra, S., Srivastava, R.C., et al. (2022). Impact of soil-water contaminants on tropical agriculture, animal, and societal environment. *Advances in Agronomy*, 176:209–274.

²² Abdelmotelb, K.F., Prasad, S.H., Lavale, S.A., Chichaghare, A.R. & Mehta, S. (2023). Radioisotopes and their impact on plants. In: *Plants and Their Interaction to Environmental Pollution*. Elsevier; 2023. P. 283–98.

3.2.5 Biomedical and Pharmaceutical Waste

The COVID-19 pandemic led to an exponential increase in biomedical waste, especially single-use plastics, antiviral drugs, and disinfectants. These materials often contain polymeric additives, microplastics, and residual pharmaceuticals that do not degrade easily. When dumped on land, they not only introduce infectious risks but also promote antibiotic resistance in soil microbiota²³. This has broad implications for public health, given the global rise in drug-resistant pathogens.

3.3 Modeling Pollutant Mobility and Persistence

Understanding the transport and retention of hazardous substances in soil can be approached through the following generalized contaminant transport equation:

$$\frac{\partial C}{\partial t} = D \frac{\partial^2 C}{\partial x^2} - v \frac{\partial C}{\partial x} - \mu C \quad (2)$$

Where:

- C = contaminant concentration (mg/L),
- D = dispersion coefficient (cm²/day),
- v = average pore water velocity (cm/day),
- μ = degradation or sorption rate constant (day⁻¹),
- x = spatial coordinate (cm),
- t = time (days).

This one-dimensional advection-dispersion-reaction equation captures how a contaminant spreads, moves with groundwater, and is attenuated *via* chemical reactions or microbial degradation. For heavy metals, the value of μ is often near zero, reflecting their persistence, whereas for biodegradable organics, μ may vary significantly with microbial abundance and soil redox conditions²⁴. Figure 1 shows how hazardous waste enters the soil through surface or subsurface deposition. Contaminants bind to particles, infiltrate to groundwater, or are taken up by vegetation. Biota and microbial processes are disrupted, leading to ecosystem imbalance.

²³ Zhu, Y. G., Zhao, Y., Zhu, D., Gillings, M., Penuelas, J., Ok, Y. S., Capon, A., & Banwart, S. (2019). Soil biota, antimicrobial resistance and planetary health. *Environment International*, 131, 105059. <https://doi.org/10.1016/j.envint.2019.105059> | Book series

²⁴ Padhye, L. P., Srivastava, P., Jasemizad, T., Bolan, S., Hou, D., Shaheen, S. M., Rinklebe, J., O'Connor, D., Lamb, D., Wang, H., Siddique, K. H. M., & Bolan, N. (2023). Contaminant containment for sustainable remediation of persistent contaminants in soil and groundwater. *Journal of Hazardous Materials*, 455, 131575. <https://doi.org/10.1016/j.jhazmat.2023.131575>

3.4 Visualizing the Impact




Figure 1: Conceptual model of hazardous waste interactions with soil systems

4. Innovative Remediation Technologies: A Transformational Shift

Contaminated land poses an escalating risk to environmental and human health, and while the urgency to address this issue has been acknowledged for decades, remediation approaches have traditionally relied on “dig-and-dump” methods. Excavation and offsite landfilling—while effective for immediate containment—are energy-intensive, prohibitively costly, and risk merely transferring contamination to another location without truly eliminating the hazard²⁵.

In contrast, the scientific community is now embracing a paradigm shift toward sustainable, in situ remediation technologies—solutions that act within the contaminated site, reducing ecological disruption and long-term costs. These emerging tools integrate advancements in biotechnology, nanotechnology, electrochemistry, and materials science. What follows is an in-depth exploration of five such technologies that represent the frontier of land decontamination.

4.1 Phytoremediation with Gene-edited Plants

Phytoremediation uses the natural uptake, accumulation, or degradation capacity of plants to remove contaminants from soil. Hyperaccumulators such as *Brassica juncea* (Indian mustard) and *Vetiveria zizanioides* (vetiver grass) have demonstrated strong potential to sequester heavy metals

²⁵ Jain, H. (2024). From pollution to progress: Groundbreaking advances in clean technology unveiled. *Innovation and Green Development*;3(2):100143.

like cadmium, lead, and arsenic²⁶. However, conventional phytoremediation is often slow and limited by biomass and root depth.

After the emergence of CRISPR-Cas9 gene editing, scientists are able to engineer hyperaccumulators to make them have maximum exudation of roots, metal transporters, and degradation enzymes. To give an example, metal transporter encoding genes (especially AtHMA4 and PCS1) are highly overexpressed in an effort to enhance cadmium uptake in *Arabidopsis thaliana*²⁷. Moreover, engineering of plant-specific bacterial symbionts into a bioengineered rhizosphere system produces greater specificity and toleration to pollutants of the plant-microbe system.




Figure 2: CRISPR-enhanced phytoremediation system for heavy metal uptake

4.2 Nano-enabled Soil Remediation

Nanotechnology brings about a new set of materials that can interact with soil contaminants at a molecular level. As an example, nanoscale zero-valent iron (nZVI) presents an outstanding reactivity as a result of the large surface area and the ability to donate electrons. It has worked well with respect to the immobilization of arsenic, reduction of hexavalent chromium, as well as degradation of chlorinated solvents²⁸.

The remediation mechanism can be expressed by the first-order degradation kinetics:

$$\frac{dC}{dt} = -kC \quad (3)$$

²⁶ Otunola, B.O., Aghoghovia, M.P., Thwala, M. & Ololade, O.O. (20220). Heavy metal phytoremediation potential of Vetiver grass and Indian mustard: update on enhancements and research opportunities. *Water Air Soil Pollution*. 2022;233(5):154.

²⁷ Nosek, M., Kaczmarczyk, A., Jędrzejczyk, R.J., Supel, P., Kaszycki, P., Miszański, Z. (2020). Expression of genes involved in heavy metal trafficking in plants exposed to salinity stress and elevated Cd concentrations. *Plants*;9(4):475.

²⁸ Sun, X., Fang, J.M., Xu, R., Wang, M., Yang, H., Han, Z., et al. (2024). Nanoscale zero-valent iron immobilized inside olidification/stabilization. *Journal of Environmental Chemical Engineering*; 12(5):113846.

Where:

- C is the contaminant concentration,
- k is the rate constant (day^{-1}),
- t is time.

The high k value associated with nZVI reflects its rapid contaminant attenuation, especially under anaerobic conditions. Moreover, nanoparticles such as TiO_2 and ZnO exhibit photocatalytic properties, enabling degradation of persistent organic pollutants under UV exposure²⁹. However, caution must be exercised as the environmental fate and ecotoxicology of free nanoparticles remain under scrutiny. Innovations like bio-nano conjugates and encapsulated nanomaterials are being explored to mitigate off-target effects.

4.3 Electrokinetic Remediation (EKR)

Electrokinetic remediation is particularly effective in fine-grained soils (like clay), where traditional fluid-based extraction methods fail due to low permeability. EKR applies a low-voltage direct current across electrodes embedded in the soil, inducing the migration of charged contaminants through electroosmosis, electrophoresis, and electromigration³⁰. EKR systems can be enhanced using chelating agents (e.g., EDTA or citric acid), surfactants, or even microbial consortia, thereby enabling the removal of heavy metals, hydrocarbons, and radionuclides. The electrokinetic flux J is defined by:

$$J = \mu_e EC \quad (4)$$

Where:

- μ_e is electrophoretic mobility ($\text{cm}^2/\text{V}\cdot\text{s}$),
- E is electric field strength (V/cm),
- C is the contaminant concentration.

²⁹ Musa, I.O., Ijah, U.J.J., Abioye, O.P., Abdulsalam, M., Pal, S. K., Abamhekelu, I.A., et al. (2024). Phytoremediation of Heavy Metal-Contaminated Soil. In: Eco-Restoration of Polluted Environment. CRC Press; 2024. P. 35–52.

³⁰ Fanali, C., Della Posta, S., Gentili, A., Chankvetadze, B., & Fanali, S. (2023). Recent developments in electromigration techniques related to pharmaceutical and biomedical analysis - A review. *Journal of Pharmaceutical and Biomedical Analysis*, 235, 115647. <https://doi.org/10.1016/j.jpba.2023.115647>.




Figure 3: Conceptual diagram of electrokinetic remediation mechanism

4.4 Microbial Consortia and Synthetic Biology


Bioremediation has entered a new phase, powered by synthetic biology. Traditional single-strain bacterial treatments are now being replaced by engineered consortia that operate collaboratively under quorum-sensing frameworks to degrade complex mixtures of pollutants. Another example is that microbial consortia (*Pseudomonas putida*, *Deinococcus radiodurans*, and *Rhodococcus erythropolis*) are modified synthetically to not only degrade hydrocarbons but also heavy metals at the same time³¹. These organisms can be made to respond to a given chemical signal by the addition of synthetic gene circuits, hence reducing off-target activity. What is more, degradation rates may be increased by bioaugmentation with CRISPR-edited strains without a significant impact on the native microbial diversity. The difficulty lies in the control of their release and their survival within natural environments, and the creation of self-limiting gene circuits and kill-switch technologies is underway.

4.5 Geopolymerization

Geopolymerization represents a sustainable route for the immobilization of hazardous waste within a stable, inert matrix. Derived from industrial by-products like fly ash, slag, and metakaolin, geopolymers are inorganic aluminosilicate polymers that form a three-dimensional lattice structure under alkaline activation.

³¹ Qin R, Xu T, Jia X. (2022). Engineering *Pseudomonas putida* to produce rhamnolipid biosurfactants for promoting phenanthrene biodegradation by a two-species microbial consortium. *Microbiology Spectrum*; 10(4): e00910-22.

The chemical process follows the reaction:



This reaction immobilizes contaminants by physically encapsulating them and chemically binding to metal ions. Studies have shown that geopolymers can reduce the leachability of lead, cadmium, and arsenic by more than 95% ³².

5. Interdisciplinary Pathways for Sustainable Remediation

The remediation of hazardous land-based waste is not merely a scientific or technological endeavor; it is a multifaceted societal challenge that spans ecology, engineering, data science, public policy, and ethics. While recent innovations in remediation have demonstrated promise, the complex and dynamic nature of contaminated landscapes requires more than isolated technical solutions. No single method, phytoremediation, nanotechnology, or electrochemical treatment—can universally address the diverse array of contaminants or the sociopolitical contexts in which they exist ³³. What is increasingly evident is that sustainable remediation depends on interdisciplinary integration: bringing together the precision of environmental monitoring, the power of computational modeling, the inclusivity of participatory research, and the legitimacy of policy co-design. This convergence of science, community, and governance offers a pathway to interventions that are not only effective, but also equitable and enduring.

Table 3. Comparative summary of innovative remediation technologies.

Technology	Target Contaminants	Key Advantage	Key Limitation
Gene-edited Phytoremediation	Heavy metals, organics	Low-cost, solar-powered	Slow rate; depth-limited
Nano-remediation	Metals, solvents, and POPs	High reactivity, fast kinetics	Potential nanoparticle toxicity
Electrokinetic Remediation	Ions, metals, and hydrocarbons	Works in low-permeability soils	Energy consumption
Synthetic Microbial Consortia	Hydrocarbons, heavy metals, and pharmaceuticals	High specificity, multi-contaminant capacity	Regulatory and ecological concerns
Geopolymerization	Inorganics (heavy metals, ash)	Long-term immobilization, reuse of waste	Not suitable for organic pollutants

³² Pu, S., Shen, Z., Duan, W., Lang, L., Liu, Y., Xu, B., et al. (2021). Discussion on the applicability and mechanism of phosphate-based geopolymers used for cadmium and cadmium-lead heavy metals s engineers *Eisenia fetida*. *Ecotoxicology and Environmental Safety*; 227:112891.

³³ Phang, L. Y., Mingyuan, L., Mohammadi, M., Tee, C. S., Yuswan, M. H., Cheng, W. H., & Lai, K. S. (2024). Phytoremediation as a viable ecological and socioeconomic management strategy. *Environmental Science and Pollution Research International*, 31(38), 50126–50141. <https://doi.org/10.1007/s11356-024-34585-z>

5.1 Smart Sensor Networks for Real-time Monitoring

Contamination in soil is often invisible, spatially heterogeneous, and temporally dynamic. Traditional laboratory-based sampling is time-consuming, spatially limited, and often fails to capture real-time fluctuations. The deployment of smart sensor networks—based on Internet of Things (IoT) architecture enables real-time, in situ monitoring of key soil parameters such as heavy metal concentration, redox potential, temperature, and pH³⁴. For instance, electrochemical sensors embedded with nanomaterials can detect trace levels of cadmium or arsenic at the parts-per-billion (ppb) scale, offering early warning capabilities for leaching events. Wireless sensor arrays can then relay this information through cloud-based platforms for spatial analysis and response.




Figure 4: Framework of smart sensor network integration for land contamination monitoring

5.2 Machine Learning Models for Predictive Remediation

Contaminant fate and transport in soil systems are governed by non-linear interactions involving hydrology, soil structure, microbial ecology, and chemical kinetics. Traditional models such as the advection-dispersion equation struggle to capture such complexity. Machine learning (ML) and artificial intelligence (AI) offer the capability to analyze high-dimensional data from sensor networks, historical contamination records, and soil property databases. Algorithms such as Random Forests, Long Short-Term Memory (LSTM) neural networks, and Gaussian Process Regression can now predict contaminant plumes, identify hotspots, and simulate the efficacy of different remediation strategies under varying environmental conditions³⁵. For example, a recent LSTM model trained on over 10 years of soil cadmium data from East China achieved a root mean

³⁴ Gao, J., Han, H., Gao, C., Wang, Y., Dong, B., & Xu, Z. (2023). Organic amendments for in situ immobilization of heavy metals in soil: A review. *Chemosphere*, 335, 139088. <https://doi.org/10.1016/j.chemosphere.2023.139088>

³⁵ Saheed, Y.K., Salau-Ibrahim, T.T., Abdulsalam, M., Adeniji, I.A. & Balogun, B.F. (2024) Modified bi-directional long short-term memory and hyperparameter tuning of supervised machine learning models for cardiovascular heart disease prediction in the mobile cloud environment. *Biomed Signal Process Control*. 2024;94:106319.

square error (RMSE) of 0.21 mg/kg, outperforming conventional models in forecasting pollutant migration³⁶.

Mathematically, predictive ML models minimize the loss function:

$$L(y, \hat{y}) = \frac{1}{n} \sum_{i=1}^n (y_i, \hat{y}_i)^2 \quad (5)$$

where:

- y_i = actual contaminant concentration,
- $\hat{y}_i$ = predicted concentration,
- n = number of samples.

Such models support adaptive management by simulating outcomes before field interventions are launched.

5.3 Policy–Technology Co-design

For remediation to be implementable on a scale, it must be socially acceptable, politically feasible, and economically viable. Too often, technological interventions are introduced without engaging the communities they intend to serve, leading to mistrust, abandonment, or conflict. Policy–technology co-design addresses this by involving multi-stakeholder actors—including scientists, policymakers, NGOs, and affected residents—from the earliest stages of research and development³⁷. An example of successful co-design is seen in Brazil's contaminated urban peripheries, where city councils collaborated with environmental engineers and slum communities to develop low-cost bioremediation systems using native plants and organic composting³⁸. These efforts improved soil quality while building public trust in science.

5.4 Community-based Participatory Research (CBPR)

Beyond co-design, community-based participatory research empowers residents to become co-researchers in environmental assessment and remediation. CBPR initiatives have been particularly effective in Indigenous and historically marginalized communities, where legacies of exploitation and distrust of scientific institutions are prevalent. In New Mexico, for example, Navajo communities have partnered with university scientists to monitor uranium contamination from abandoned mines, using

³⁶ Zhao, W., Ma, J., Liu, Q., Dou, L., Qu, Y., Shi, H., Sun, Y., Chen, H., Tian, Y., & Wu, F. (2023). Accurate Prediction of Soil Heavy Metal Pollution Using an Improved Machine Learning Method: A Case Study in the Pearl River Delta, China. *Environmental Science & Technology*, 57(46), 17751–17761. <https://doi.org/10.1021/acs.est.2c07561>.

³⁷ Mugari E, Nethengwe NS, Gumbo AD. (2025). A co-design approach for stakeholder engagement and knowledge integration in flood risk management in Vhembe district, South Africa. *Front Clim*. 2025;7:1517837.

³⁸ Shrivastava NG. (2021). Bioremediation of solid waste management. In: *Handbook of Solid Waste Management: Sustainability Through Circular Economy*. 2021. P. 1–51.

locally collected samples and culturally adapted educational tools ³⁹. This approach builds local capacity and ensures that remediation addresses not just chemical hazards, but cultural, spiritual, and historical dimensions of land.

5.5 Integration is Not Optional

The effectiveness of future remediation strategies will be measured not only by how well they remove pollutants but by how well they incorporate justice, resilience, and local knowledge.

Figure 5 summarizes the interdisciplinary model required for holistic remediation.




Figure 5: Interdisciplinary framework for sustainable hazardous waste remediation

6. Future Directions and Policy Implications

The twenty-first century's environmental legacy will be defined, in part, by how we choose to confront the growing crisis of hazardous waste contamination. While recent advances in biotechnology, nanomaterials, and systems engineering have ushered in powerful tools for site remediation, technical innovations alone are insufficient to reverse decades of degradation or prevent future harm. A sustainable path forward demands a fundamental realignment of science, policy, and societal priorities. Hazardous waste is not just a byproduct of industrial processes; it is a systemic manifestation of unsustainable economic models, regulatory fragmentation, and environmental injustice. As such, the response must be equally systemic—bridging the divide between environmental science and socio-political reform.

6.1 Enforcing the Cradle-to-Grave Principle

A central policy priority is the enforcement of cradle-to-grave tracking—a regulatory framework that mandates accountability for hazardous materials from production through disposal. While instruments such as the Basel Convention have attempted to address transboundary movements of hazardous waste, national-level enforcement remains uneven and underfunded.

In high-income countries, digitized waste tracking systems—using blockchain or geospatial monitoring—have demonstrated success in minimizing illegal dumping and enhancing

³⁹ Rock, T., & Ingram, J. C. (2020). Traditional Ecological Knowledge Policy Considerations for Abandoned Uranium Mines on Navajo Nation. *Human Biology*, 92(1), 19–26. <https://doi.org/10.13110/humanbiology.92.1.01>

compliance⁴⁰. In contrast, many low- and middle-income nations lack the institutional capacity to track hazardous materials beyond industrial gates. Investments in waste infrastructure, transparent registries, and international technical cooperation are urgently needed to close this governance gap.

6.2 Public and Private Investment in Green Remediation R&D

Governments must play a catalytic role by incentivizing research and development in low-carbon, low-impact remediation technologies. Green remediation not only emphasizes energy efficiency and reduced emissions during cleanup but also aligns with broader goals of the circular economy and land restoration. This can include subsidies for startup incubators focused on bioremediation, tax incentives for industries that adopt waste valorization techniques, and public-private partnerships to scale promising field technologies⁴¹. Academic institutions should also be supported in establishing interdisciplinary remediation laboratories that merge environmental engineering, microbiology, and materials science.

6.3 Scientific Gaps and Research Needs

From a scientific standpoint, the remediation field must evolve to address persistent gaps that limit translatability and scalability.

Field-scale Validation Over Lab Reliance

Many remediation technologies perform well in controlled laboratory conditions but falter in heterogeneous, real-world settings. Variables such as fluctuating temperature, soil heterogeneity, and co-contaminant interference often reduce efficacy. There is a pressing need for large-scale field trials across different climatic zones and soil types, with independent monitoring and open-access datasets⁴².

Co-contaminant Interactions

Most contaminated sites are exposed to complex mixtures of pollutants, for example, metals coexisting with organic solvents or radionuclides. However, studies tend to focus on single-contaminant systems, overlooking interactions that may alter mobility, toxicity, or degradability⁴³. Future research should incorporate multi-factorial designs that mirror environmental realities, supported by mechanistic modeling of synergistic or antagonistic effects.

Longitudinal Soil Microbiome Studies

⁴⁰ D'Amato A, Mazzanti M, Nicolli F, Zoli M. (2018). Illegal waste disposal: Enforcement actions and decentralized environmental policy. *Socio-Economic Planning Sciences*;64:56–65.

⁴¹ Imarhiagbe, O., Nwodo, M.U., Ogwu, M.C. (2024). Innovations for Cleaner Futures: Emerging Technologies in Air Pollution Mitigation. In: *Sustainable Strategies for Air Pollution Mitigation*. 2024. P. 377–407.

⁴² Vijay, V., Shreedhar, S., Adlak, K., Payyanad, S., Sreedharan, V., Gopi, G., Sophia van der Voort, T., Malarvizhi, P., Yi, S., Gebert, J. & Aravind, P. (2021) Review of Large-Scale Biochar Field-Trials for Soil Amendment and the Observed Influences on Crop Yield Variations. *Frontiers in Energy Research*, 9:710766. doi: 10.3389/fenrg.2021.710766

⁴³ Innocent, M.O., Patrick, O.I., Job, O.S., Abdulsalam, M., Adams, M., Sikirula, A., et al. (2024). Introduction to Nanotoxicology. In: *Environmental Nanotoxicology: Combatting the Minute Contaminants*. Cham: Springer Nature Switzerland; 2024. P. 1–22.

Soil health is intricately tied to microbial diversity, yet the long-term recovery trajectories of microbial community's post-remediation remain poorly understood. Emerging sequencing technologies, such as metagenomics and metabolomics, can provide insight into how microbial functions recover or shift after contamination events and cleanup efforts ⁴⁴. Understanding these dynamics is vital not just for ecological restoration but also for evaluating unintended consequences of remediation interventions that may sterilize or restructure microbial networks critical for nutrient cycling and plant health.

Standardized Lifecycle Assessments (LCA)

The absence of a globally accepted system of analyzing the lifecycle of remediation technologies makes it challenging to evaluate and make a comparative analysis of their sustainability in a fair way. The existing lifecycle assessment (LCA) tools show a broad and varying range of scope and methodology, contributing to inconsistent energy consumption, carbon emission, and residual toxicity evaluation ⁴⁵. This could be achieved by setting up a uniform, peer-reviewed system of how LCAs in remediation should be conducted, which would help give policymakers a clearer way forward to prioritize their focus on technologies that fit within both the climate and population health agendas.

6.4 Human and Institutional Dimension

Technical solutions are not the only methods of effective remediation, but rather the involvement of the people and their trust and backing of the institutions. Policies should require active engagement of local communities in the process of reclaiming the hazardous land, exchange of information about risk and benefits in a transparent manner and protect the entitlements of disproportionately impacted groups. It is possible to use such frameworks as the U.S. EPA 2023 Environmental Justice (EJ) Action Plan or the Green Deal roadmap of the European Union to define the way of equitable remediation. On the international scale, organizations such as UNEP, the World Health Organization, and the World Bank are well-placed to assist with the much-needed financial and technical support that most nations require in curbing long-term pollution issues.

7. Conclusion

Soil is far more than a passive substrate; it is a living, breathing system—an ecological membrane that anchors biodiversity, sustains food production, and regulates global biogeochemical cycles. Yet, as industrial expansion accelerates and consumption intensifies, land has increasingly become a silent repository for hazardous waste: a burden it was never designed to bear. This study has highlighted how hazardous land-based waste—from heavy metals and persistent organic pollutants to biomedical residues and radionuclides—poses complex threats not only to soil health but to broader human and planetary systems. These contaminants do not operate in isolation; they ripple outward through food webs, water cycles, and social structures, often disproportionately harming

⁴⁴ Nam, N.N/, Do, H.D.K., Loan, Trinh, K.T.& Lee, N.Y. (2023). Metagenomics: An effective approach for exploring microbial diversity and functions. *Foods*; 12(11):2140.

⁴⁵ Mirabella, N., Allacker, K. & Sala, S. (2019). Current trends and limitations of life cycle assessment applied to the urban scale: critical analysis and review of selected literature. *The International Journal of Life Cycle Assessment*; 24:1174–93.

the most vulnerable communities. The resulting landscape is not only chemically compromised but ethically troubling.

As humanity navigates the precarious terrain of the Anthropocene—marked by climate instability, biodiversity loss, and resource scarcity, the urgency of confronting hazardous waste takes on a new dimension. It is no longer sufficient to treat contamination as a technical or regulatory issue alone. Rather, how we choose to address land-based hazardous waste will serve as a defining measure of our scientific maturity, societal priorities, and ethical commitments. The innovative remediation technologies published in this study, such as gene-edited phytoremediation, nano-enabled catalysis, electrokinetic treatments, and synthetic biology, provide a radical set of solutions. But what is holding promise is integration. The future does not only require innovation, but also convergence: convergence of disciplines, institutions, and communities. Such convergence is the only way in which extractive land use can be transformed into regenerative.

In addition to this, remediation should also be redefined as a three-dimensional, inclusive process, where the focus shifts more towards justice than readiness and efficiency, and where the cultural, historical, and spiritual aspects of land must be honored. It has to focus not only on chemical clean-up but ecological recovery and social renewal. Cleaning up of hazardous waste sites is consequently not simply an issue of environmental management; it is an ethical concern, an examination of our willingness to remedy the environmental damage that we have caused and to leave the earth in a better condition for our posterity. In reclaiming contaminated soils, we also reclaim a deeper human responsibility: to steward the Earth with care, humility, and foresight. The decisions we make today regarding hazardous waste will echo for centuries in the fertility of our soils, the safety of our food, and the health of our children. It is within our collective power—and indeed our obligation—to ensure that the land remains a source of life, not a legacy of harm.